

The essence of entanglement

Časlav Brukner¹, Marek Żukowski² and Anton Zeilinger¹

¹*Institut für Experimentalphysik, Universität Wien, Boltzmanngasse 5, 1090 Wien, Austria*

²*Instytut Fizyki Teoretycznej i Astrofizyki Uniwersytet Gdański, PL-80-952 Gdańsk, Poland*

Entanglement, according to Erwin Schrödinger *the essence* of quantum mechanics, is at the heart of the Einstein-Podolsky-Rosen paradox and of the so called quantum-nonlocality – the fact that a local realistic explanation of quantum mechanics is not possible as quantitatively expressed by violation of Bell's inequalities. Even as entanglement gains increasing importance in most quantum information processing protocols, its conceptual foundation is still widely debated. Among the open questions are: What is the conceptual meaning of quantum entanglement? What are the most general constraints imposed by local realism? Which general quantum states violate these constraints? Developing Schrödinger's ideas in an information-theoretic context we suggest that a natural understanding of quantum entanglement results when one accepts (1) that the amount of information per elementary system is finite and (2) that the information in a composite system resides more in the correlations than in properties of individuals. The quantitative formulation of these ideas leads to a rather natural criterion of quantum entanglement. Independently, extending Bell's original ideas, we obtain a single general Bell inequality that summarizes all possible constraints imposed by local realism on the correlations for a multi-particle system. Violation of the general Bell inequality results in an independent general criterion for quantum entanglement. Most importantly, the two criteria agree in essence, though the two approaches are conceptually very different. This concurrence strongly supports the information-theoretic interpretation of quantum entanglement and of quantum physics in general.

CONTENTS

I. INTRODUCTION	1
II. FINITENESS OF INFORMATION	2
III. INFORMATION CONTAINED IN CORRELATIONS	3
IV. QUANTIFYING INFORMATION	4
V. ALL BELL INEQUALITIES FOR CORRELATIONS	5
VI. N QUBITS THAT VIOLATE LOCAL REALISM	7
VII. CONCLUSIONS	9
ACKNOWLEDGEMENTS	9
References	9

I. INTRODUCTION

In their seminal paper in 1935 Einstein, Podolsky and Rosen (EPR) consider quantum systems consisting of two particles such that, while neither position nor momentum of either particle is well defined, both the sum of their positions and the difference of their momenta are both precisely defined. It then follows that measurement on either position or momentum performed on, say, particle 1 immediately implies for particle 2 a precise position or momentum respectively even when the two particles

are separated by arbitrary distances without any actual interaction between them.

Motivated by EPR Schrödinger (1935a) in his paper entitled "The present situation in quantum mechanics" wrote succinctly "Maximal knowledge of a total system does not necessarily include total knowledge of all its parts, not even when these are fully separated from each other and at the moment are not influencing each other at all" and he coined the term¹ "entanglement of our knowledge" to describe this situation. A central goal of our work is to take Schrödinger's position as a starting point for a quantitative information-theoretic definition of entanglement. Therefore, carefully reading his sentence one may identify three independent ideas on which Schrödinger builds his notion of entanglement. (a) First, our knowledge, or total information, of a system is bound. (b) Second, the total information of a composite system is not necessarily fully contained in its individual constituents. (c) And the third, these statements are independent of the relative space-time arrangements of the individual observations on the constituents of the composite system. Furthermore we underline that Schrödinger talks about quantum states representing our expectation catalogs and he carefully avoids the notion of properties of systems.

In 1964 John Bell obtained certain bounds (Bell inequalities) on combinations of statistical correlations for measurements on two-particle systems if these correlations are understood within a realistic picture based on local hidden properties of each individual particle. In a realistic picture the measurement results are determined by properties the particles carry prior to and independent of observation. In a local picture the results obtained at one location are independent of any measurements or actions performed at space-like separation. Then Bell showed that quantum mechanics predicts violation of these constraints for certain statistical predictions for two-particle systems. A more striking conflict between quantum mechanical and local realistic predictions even for perfect correlations has been discovered for three and more particles (Greenberger *et al.*, 1989; 1990; Mermin, 1990), resulting in an outright invalidation of the EPR

¹Schrödinger (1935a) used the German original "Verschränkung" and he himself introduced the English translation "entanglement" (Schrödinger, 1935b).

concepts. By now a number of experiments (Freedman and Clauser, 1972; Aspect *et al.*, 1981; Pan *et al.*, 2000) have confirmed the quantum mechanical predictions even if the individual particles are truly space-like separated (Weihs *et al.*, 1998). However two important questions remain: (1) "What are the most general constraints on correlations imposed by local realism?" and (2) "Which quantum states violate these constraints?". The latter has been solved in general only in the case of two particles in pure states (Gisin, 1991; Gisin and Peres 1992) and for two-qubit mixed states (Horodecki *et al.*, 1995). Only recently bounds for local realistic description of higher-dimensional systems have been found in some simple cases (Kaszlikowski *et al.*, 2000).

Our paper will now develop the ideas presented above in various quantitative ways. Following first Schrödinger's ideas we will define quantum entanglement as a feature of a composite system to have more information contained in correlations than any classical mixture of its individual constituents could ever have. The essence of classical correlations is that there the joint properties can be reduced to correlations between properties of the individual constituents. This is an operational definition because information is always defined through observation of measurement results.

In parallel, following then Bell's ideas, we will obtain a single general Bell inequality that summarizes all possible local realistic constraints on the correlations for a multi-qubit system, where two dichotomic observables are measured on each individual qubit. This enables us to introduce another operational definition of entangled states as those which violate that general Bell inequality in a direct measurement². Finally we show that the two operational definitions are equivalent in the two-qubit case and that in general our informational definition of entanglement provides a necessary and rather stringent condition for violating the general Bell inequality. We find this intriguing because the two operational approaches are based on completely different concepts. To us this further supports the view that information is the most fundamental concept in quantum physics (Zeilinger, 1999).

²By an entangled state is often meant a state which cannot be represented as a classical mixture of product states (non-separable state). Here, since our aim is to relate Schrödinger's notion of entanglement to the one via Bell's inequality we will use the definition as given in the text above. However, it is well known that there are cases of "hidden non-locality" where a quantum state initially does not directly violate a Bell inequality but after local operations together with classical communication such a violation might occur (Popescu, 1995).

II. FINITENESS OF INFORMATION

A central point in our discussion will be the different ways how information can be distributed within a composite system. We therefore have to introduce our notion of information and we have to give our considerations about how much information altogether a system can represent.

Any physical description of a physical system is a set of propositions together with their truth values - true or false. Then, any proposition we might assign to a quantum system which is always based on observation of properties of the classical apparatus used, represents our knowledge, i.e., information, of a system gained through observation. To illustrate this consider the state $|\psi\rangle = |z+\rangle$ of a spin-1/2 particle with spin up along the z -axis, which is an eigenstate of the operator σ_z with eigenvalue $+1$. This simply means that the quantum system described by the state $|\psi\rangle$ will be found with certainty to have spin $+1$ if it is measured along the z -axis. Thus the information content in that state is represented by the truth value of the proposition: "The spin along the z -axis is up." This is one bit of information. It is clear that both truth values must be possible. Only then observation of the system can result in a gain of information. In agreement with Schrödinger's idea (a) above the most simple system then just represents one bit of information. This is what we mean when we talk about a system carrying information. To us a system is a construct based on information.

If we assume that one bit is the only information the most simple quantum system can carry, and that this is defined with respect to a certain measurement then other measurements must contain an element of irreducible randomness. Otherwise the system would carry more information in conflict with (a) for a system representing one bit of information only. This means that there are other measurement directions for which the experimental outcome is completely random. Specifically, for a measurement along any direction $\vec{n}$ in the x - y plane the proposition "The spin along the $\vec{n}$ axis is up" is completely indefinite, i.e., we have absolutely no knowledge which particular outcome "spin up" or "spin down" will be observed in an individual experimental trial. The two propositions about spin along z and about spin along $\vec{n}$ are propositions with a property of mutual exclusiveness. This is quantum complementarity: the complete knowledge of the truth value of one of the propositions implies maximal uncertainty about the truth values of the other.

How much information is carried by a system with respect to a specific set of mutually complementary propositions? We suggest that it is natural to assume that the information contained in a set of mutually complementary propositions is the sum over the measures of information of the individual members of that particular set

(Bohr, 1958). Specifically, to obtain the total information carried by a quantum system one summarizes over all individual measures of information for a complete set of mutually complementary measurements, as shown in Brukner and Zeilinger (1999). There it was shown that the total information carried by the composite system consisting of N qubits in a pure state to be N bits of information.

Here we are interested in the various ways how information can be distributed within a composite system. In particular we will consider that part of the total information of the system which is contained in correlations, or joint properties of its constituents. This is also the reason why here we do not consider complete sets of mutually complementary propositions for the composite system but just a subset of them concerning joint properties of its constituents.

III. INFORMATION CONTAINED IN CORRELATIONS

Correlations between quantum systems have assumed a very central role in the discussions of the foundations of quantum mechanics. We will now investigate how much information can be contained in such correlations in order to give an information-theoretic criterion of quantum entanglement. As it is our final goal to compare that criterion with the one given by Bell-type inequalities (Clauser *et al.*, 1969; Mermin, 1990; Belinskii and Klyshko, 1993) where one considers correlations between spin measurements confined on each side within one plane we restrict our analysis to an x - y plane locally defined for each subsystem.

For clarity of presentation we first investigate the case of a two-qubit system carrying therefore $N=2$ bits of information, i.e. representing the truth value of two propositions. That information contained in 2 propositions can be distributed over the 2 qubits in various ways.

Consider first a product state e.g. $|\psi\rangle = | +x\rangle_1 | -x\rangle_2$. Here the state $|\psi\rangle$ represents the two-bit combination true-false of the truth values of the propositions about the spin of each particle along the x -axis: (1) "The spin of particle 1 is up along x " and (2) "The spin of particle 2 is up along x ". Instead of the second proposition describing the spin of particle 2, we could alternatively choose a proposition which describes the result of a joint observation: (3) "The two spins are the same along x ." Then the state $|\psi\rangle$ represents the two-bit combination true-false of the truth values of the propositions (1) and (3). Note that this is also present in classical composite systems.

Evidently, for pure product states at most one proposition with definite truth-value can be made about joint properties because one proposition has to be used up to define a property of one of the two subsystems. In other words 1 bit of information defines the correlations

$$I_{corr}^{prod} = 1. \quad (1)$$

In our example where $|\psi\rangle = | +x\rangle_1 | -x\rangle_2$ the correlations are fully represented by the correlations between x -measurements on both sides, therefore $I_{corr}^{prod} = 1 = I_{xx} = 1$. The specific measure of information used by us will be specified below.

Obviously, the choice of directions within the planes of measurement on the two sides is arbitrary. Choosing general x and y directions we request that the total information contained in the correlations must be invariant upon this choice. This invariance property by itself already defines the specific measure of information and it rules out Shannon's measure (Brukner and Zeilinger, 2001). Following our arguments given above we now define the information contained in the correlations as the sum over the individual measures of information carried in a complete set of mutually complementary observations within x - y plane. Therefore the information contained in the correlations is quantified by the sum

$$I_{corr} = I_{xx} + I_{xy} + I_{yx} + I_{yy} \quad (2)$$

of the partial measures of information contained in the set of complementary observations within the x - y -plane. These observations are mutually complementary for *product states* and the set is complete as there exists no further complementary observation within the chosen planes. By this we mean that for any product state a complete knowledge contained in one of the observations in Eq. (2) excludes any knowledge content in the other three observations.

Consider now a maximally entangled Bell state, e.g.

$$\begin{aligned} |\phi^-\rangle &= \frac{1}{\sqrt{2}}(| +x\rangle_1 | -x\rangle_2 + | -x\rangle_1 | +x\rangle_2) \\ &= \frac{1}{\sqrt{2}}(| +y\rangle_1 | +y\rangle_2 + | -y\rangle_1 | -y\rangle_2). \end{aligned} \quad (3)$$

The two propositions here both are statements about results of joint observations (Zeilinger 1997), namely (1') "The two spins are equal along x " and (2') "The two spins are equal along y ". Now the state represents the two-bit combination false-true of these propositions about correlations. Note that here the 2 bits of information are all carried by the 2 qubits in a joint way, with no individual qubit carrying any information on its own. In other words, as the two available bits of information are already exhausted in defining joint properties, no further possibility exists to also encode information in individuals. Therefore

$$I_{corr}^{Bell} = 2. \quad (4)$$

Note that in our example $I_{xx} = I_{yy} = 1$ and $I_{xy} = I_{yx} = 0$. Also, note that the truth value for another proposition, namely, "The two spins are equal along z " must follow

immediately from the truth values of the propositions (1') and (2'), as only 2 bits of information are available. Interestingly this is also a direct consequence of the formalism of quantum mechanics as the joint eigenstate of $\sigma_x^1 \sigma_x^2$ and of $\sigma_y^1 \sigma_y^2$ is also the eigenstate of $\sigma_z^1 \sigma_z^2 = -(\sigma_x^1 \sigma_x^2)(\sigma_y^1 \sigma_y^2)$.

In contrast to product states we suggest entanglement of two qubits to be defined in general such that *more than one bit* (of the two available ones) is used to define joint properties, i.e.

$$I_{corr}^{entgld} > 1 \quad (5)$$

for at least one choice of the local x and y directions for the two qubits. This is in agreement with Schrödinger's idea (b). Equivalently we suggest to define the two qubits as classically composed (non-entangled) if less than or equal to one bit of information is used to define correlations, i.e.

$$I_{corr}^{nonent} \leq 1 \quad (6)$$

for all possible choices of the local x and y directions for two qubits. As we will show below, the independent information-theoretic definition of entanglement (5) will turn out to be equivalent to a necessary and sufficient condition for a violation of any Bell-type inequality for two-qubits.

In the generalization to more and more qubits we consider, without loss of generality, a product state $|\psi\rangle = |x_1\rangle|x_2\rangle\ldots|x_N\rangle$ of N qubits. Here, x_j denotes a spatial direction in a local coordinate system of observer j . Then only one proposition with definite truth-value can be made about the correlations in the N qubits, namely the proposition (*): "The product of spin of particle 1 along x_1 , spin of particle 2 along x_2 , ... and spin of particle N along x_N is +1". This means that for a product state or a classical mixture of product states again at most one bit is represented in N -qubit correlations. We therefore here too suggest to define N qubits as classically composed if not more than one bit of information is used to define correlations, i.e.

$$I_{corr}^{nonent} = \sum_{\substack{x_1, \dots, x_N \\ = x, y}} I_{x_1 \dots x_N} \leq 1 \quad (7)$$

for all possible choices of local x and y directions for N qubits. Here the sum is over the measures of information over a set of propositions of the type given above where $x_1, \dots, x_N \in \{x, y\}$ and which therefore are mutually complementary for product states³.

³Since the set of propositions entering Eq. (7) contains also propositions which are not mutually complementary for a non-product state the sum in (7) does not give the value of

We would like to make a very general comment. The quantitative condition (7) while certainly correct for the situations discussed here might have to be modified in order to apply to more complicated cases like entanglement between many qubits when the measurements are not restricted to one plane or entanglement between systems defined in Hilbert spaces of higher dimensions, so-called qunits. Another interesting case can arise when one also considers in detail all possible sets of correlations between all possible sets of subsystems. For example for 3-qubit systems we have one correlation between all three individual qubits, we have 3 correlations between two individual qubits and we have another 3 correlations between one qubit and the other two. This results in a large number of conditions of the type of Eq. (7) which are not independent from each other in general. Yet, we stress, that it is to be expected that the general ideas laid out here will still be applicable. A most important guidance for quantitative conditions being that the information carried by the correlations between subsystems exceeds the limit given by the information carried by the subsystems themselves.

Finally we stress that in our information-theoretic analysis of entanglement we did not have to use concepts like spatial separation between subsystems of the composite system or the relative times of the observations on the subsystems. This is in agreement with Schrödinger's idea (c).

To this point we did not yet specify any particular measure of information. Thus the question arises which particular measure of information is adequate to define the information gain in an individual quantum experiment.

IV. QUANTIFYING INFORMATION

Consider an experiment with two outcomes "yes" and "no" and with the probabilities p_1 and $p_2 = 1 - p_1$, respectively for the two outcomes. Within finite time the experimenter can perform only a finite number of experimental trials. Because of inherent fluctuations associated

information in bits contained in correlations for such states. It is then not surprising that, for example, in the case of a three-qubit GHZ state the sum results in 4 ($I_{xyy} = I_{yyx} = I_{yxx} = 1$, other are zero). Here however only three of the four propositions are independent, as only 3 bits of information are available. This again is also a direct consequence of the formalism of quantum mechanics as the joint eigenstate of $\sigma_x^1 \sigma_y^2 \sigma_y^3$, $\sigma_y^1 \sigma_x^2 \sigma_y^3$ and $\sigma_y^1 \sigma_y^2 \sigma_x^3$ is also an eigenstate of $\sigma_x^1 \sigma_x^2 \sigma_x^3 = -(\sigma_x^1 \sigma_y^2 \sigma_y^3)(\sigma_y^1 \sigma_x^2 \sigma_y^3)(\sigma_y^1 \sigma_y^2 \sigma_x^3)$. Nevertheless when the sum in (7) is larger than unity, this indicates that the total information contained in correlations is larger than one bit.

with any probabilistic experiment with a finite number of trials the number of occurrences of a specific outcome in future repetitions of the experiment is not precisely predictable. Rather it obeys the binomial distribution (See e.g. Gnedenko, 1976).

If one bets for example that the number of "yes" outcomes will be the one with highest probability, the probability of success still depends on p_1 . With a probability of $p_1 = 0.5$, the probability of 5 "yes" outcomes in 10 trials is only 0.25, but with $p_1 = 0.9$ the probability of 9 "yes" outcomes in 10 trials is 0.39. It is a trait of the binomial distribution, that we know the future number of occurrences of the outcomes very well if p_1 (or equivalently p_2) is close to 0 or 1, but we know much less about them when p_1 is around 0.5. Note that this follows from elementary probability theory without any input from physics⁴. In (Brukner and Zeilinger 1999) it was shown that this knowledge is properly represented by the measure

$$I = (p_1 - p_2)^2. \quad (8)$$

This attains its maximal value of unity when one of probabilities is one, and it attains its minimal value of 0 when both probabilities are equal.

Note that all our propositions about joint properties are binary propositions, i.e. they are associated to experiments with two possible outcomes, one of them being "the product of spin of particle 1 along x_1 , spin of particle 2 along x_2 , ... and spin of particle N along x_N is +1", and the other one "the product of spin of particle 1 along

x_1 , spin of particle 2 along x_2 , ... and spin of particle N along x_N is -1", so that Eq. (8) can be applied. If we now denote the probabilities for the two outcomes by $p_{x_1 \dots x_N}^+$ and $p_{x_1 \dots x_N}^-$ respectively, then the information contained in proposition (*) is given by

$$I_{x_1 \dots x_N} = (p_{x_1 \dots x_N}^+ - p_{x_1 \dots x_N}^-)^2. \quad (9)$$

Now we will express Eq. (9) in terms of the density matrix ρ of N qubits. First note that an arbitrary mixed state of N qubits can be written as

$$\rho = \frac{1}{2^N} \sum_{x_1, \dots, x_N=0}^3 T_{x_1 \dots x_N} \sigma_{x_1}^1 \otimes \dots \otimes \sigma_{x_N}^N \quad (10)$$

where σ_0^j is the identity operator in the Hilbert space of particle j , and $\sigma_{x_j}^j$ is a Pauli operator for $x_j = 1, 2, 3$. Here the elements of the correlation tensor T are given as mean values of the product of the N spins,

$$T_{x_1 \dots x_N} = \text{Tr}[\rho(\sigma_{x_1}^1 \otimes \dots \otimes \sigma_{x_N}^N)] = p_{x_1 \dots x_N}^+ - p_{x_1 \dots x_N}^- \quad (11)$$

with $T_{0 \dots 0} = 1$. Then obviously our measure of information (9) is equal to the square of the corresponding element of the correlation tensor

$$I_{x_1 \dots x_N} = T_{x_1 \dots x_N}^2. \quad (12)$$

We have thus obtained a quantitative expression for the individual measures of information contained in the sum of Eq. (7) and we want to emphasize that our analysis of entanglement would not be possible without the use of the measure of information (8).

So far in the present paper we followed Schrödinger's concepts of entangled states in our information-theoretic analysis. Now we will follow Bell's ideas in a second, independent approach to characterize entanglement.

V. ALL BELL INEQUALITIES FOR CORRELATIONS

Here we obtain a single general Bell inequality that summarizes all possible constraints on the statistical correlations of an N-qubit system. These constraints are derived under the assumptions of local realism. We consider such correlation measurements where for each individual particle one of the two arbitrary dichotomic observables can be chosen. From this inequality we obtain as specific corollaries the Clauser-Horne-Shimony-Holt (CHSH) inequality (Clauser *et al.*, 1969) for two-qubit systems and the related inequalities for N qubits (Mermin, 1990; Ardehali, 1992; Belinskii and Klyshko, 1993).

In a local realistic picture one assumes that the result of every measurement of an observable is predetermined. Thus in such a picture one implicitly requires an unlimited amount of information to be carried by an individual

⁴Here a very subtle position was assumed by von Weizsäcker (1975) who wrote: "It is most important to see that this [the fact that probability is not a prediction of the precise value of the relative frequency] is not a particular weakness of the objective empirical use of the concept of probability, but a feature of the objective empirical use of any quantitative concept. If you predict that some physical quantity, say a temperature, will have a certain value when measured, this prediction also means its expectation value within a statistical ensemble of measurements. The same statement applies to the empirical quantity called relative frequency. But here are two differences which are connected to each other. The first difference: In other empirical quantities the dispersion of the distribution is in most cases an independent empirical property of the distribution and can be altered by more precise measurements of other devices; in probability the dispersion is derived from the theory itself and depends on the absolute number of cases. The second difference: In other empirical quantities the discussion of their statistical distributions is done by another theory than the one to which they individually belong, namely by the general theory of probability; in probability this discussion evidently belongs to the theory of this quantity, namely of probability itself. The second difference explains the first one."

particle, which conflicts with Schrödinger's idea (a). Take therefore an individual observer and allow him or her to be able to choose between two dichotomic observables (determined by some parameters denoted here $\vec{n}_1$ and $\vec{n}_2$). This implies the existence of two numbers $A_j(\vec{n}_1)$ and $A_j(\vec{n}_2)$ each taking values $+1$ or -1 which describe the predetermined result of a measurement by the observer of the observable defined by the local parameter $\vec{n}_1$ and $\vec{n}_2$, respectively. We choose such a notation for brevity; of course each observer can choose independently two arbitrary directions.

In a specific run of the experiment the correlations between all N observations can be represented by the product $\prod_{j=1}^N A_j(\vec{n}_{k_j})$, with $k_j = 1, 2$. The correlation function is then the average over many runs of the experiment

$$E(\vec{n}_{k_1}, \dots, \vec{n}_{k_N}) = \left\langle \prod_{j=1}^N A_j(\vec{n}_{k_j}) \right\rangle_{avg}. \quad (13)$$

Note that for each observer j one has either $|A_j(\vec{n}_1) + A_j(\vec{n}_2)| = 0$ and $|A_j(\vec{n}_1) - A_j(\vec{n}_2)| = 2$ or the other way around. Then for all sign sequences of $s_1, \dots, s_N$, where $s_j \in \{-1, 1\}$ the modulus of the product $|\prod_{j=1}^N [A_j(\vec{n}_1) + s_j A_j(\vec{n}_2)]|$ vanishes except just one for which the product is 2^N . Therefore one has

$$\sum_{\substack{s_1, \dots, s_N \\ = -1, 1}} \left| \prod_{j=1}^N [s_j A_j(\vec{n}_1) + s_j^2 A_j(\vec{n}_2)] \right| = 2^N. \quad (14)$$

It then follows directly that the correlation functions must satisfy the following general Bell inequality (Weinfurter and Żukowski, 2001; Żukowski and Brukner, 2001; for an independent derivation see Werner and Wolf, 2001)

$$\sum_{\substack{s_1, \dots, s_N \\ = -1, 1}} \left| \sum_{\substack{k_1, \dots, k_N \\ = 1, 2}} s_1^{k_1} \dots s_N^{k_N} E(\vec{n}_{k_1}, \dots, \vec{n}_{k_N}) \right| \leq 2^N. \quad (15)$$

Therefore within each modulus we have sums of all 2^N correlation functions, which are the result of the Bell-type experiment, however each correlation functions is multiplied by a specific sign $\pm$. Using the generalized triangle inequality we obtain the set of Bell-type inequalities which are equivalent to inequality (15)

$$\left| \sum_{\substack{s_1, \dots, s_N \\ = -1, 1}} S(s_1, \dots, s_N) \sum_{\substack{k_1, \dots, k_N \\ = 1, 2}} s_1^{k_1} \dots s_N^{k_N} E(\vec{n}_{k_1}, \dots, \vec{n}_{k_N}) \right| \leq 2^N, \quad (16)$$

where $S(s_1, \dots, s_N)$ is one of the 2^{2^N} sign functions of $s_1, \dots, s_N$, by which we mean that its possible values can only be $+1$ or -1 . Actually, inequalities (16) represent the complete set⁵ of all possible 2^{2^N} inequalities for the

correlations, one for each possible choice of the sign function S . Many of these inequalities are trivial (for example when the choice is $S(s_1, \dots, s_N) = 1$ for all arguments, we get that the modulus of the correlation function does not exceed 1).

An exemplary stringent condition result for $S(s_1, \dots, s_N) = \sqrt{2} \cos(-\frac{\pi}{4} + (s_1 + \dots + s_N - N)\frac{\pi}{2})$ which leads to the series of inequalities derived by Belinskii and Klyshko (1993). Specifically, for $N = 2$, the CHSH inequality

$$|E(1, 1) + E(1, 2) + E(2, 1) - E(2, 2)| \leq 2 \quad (17)$$

follows. For $N = 3$, one obtains

$$|E(1, 2, 2) + E(2, 1, 2) + E(2, 2, 1) - E(1, 1, 1)| \leq 2, \quad (18)$$

where here we use numbers 1 and 2 to denote directions $\vec{n}_1$ and $\vec{n}_2$, respectively. Inequality (18) leads to the Greenberger-Horne-Zeilinger (Greenberger *et al.* 1989; 1990) contradiction for an appropriate choice of local settings. In these cases the left hand side of inequality (18) reaches the value 4 which is the maximum possible value for any, not only quantum, correlation function.

Thus far we have shown that when a local realistic model applies, the general Bell inequality (15) follows. The reverse is also true and we give the proof below: whenever inequality (15) holds one can construct a local realistic model for the correlation function. This establishes the general Bell inequality presented above as a necessary and sufficient condition for local realistic description of multi-particle correlations, where two dichotomic observables are measured on each individual particle.

The proof of the sufficiency of condition (15) will be done here in a constructive way. Simply one ascribes to the set of predetermined local results, which satisfy the following conditions $A_j(\vec{n}_1) = s_j A_j(\vec{n}_2)$, the hidden probability $p(s_1, \dots, s_N) = \frac{1}{2^N} \left| \sum_{k_1, \dots, k_N} s_1^{k_1} \dots s_N^{k_N} E(\vec{n}_{k_1}, \dots, \vec{n}_{k_N}) \right|$, and one demands that the product $\prod_{j=1}^N A_j(\vec{n}_2)$ has the same sign as that of the expression inside of the modulus defining the $p(s_1, \dots, s_N)$. In this way a definite set of local realistic values is ascribed a unique global hidden probability. Obviously, such defined probabilities are positive. However due to inequality (15) they may add up to less than 1. In such a case, the "missing" probability is ascribed to an arbitrary model of local realistic noise (e.g., for which all possible products of local results enter with equal weights). The overall contribution of such a noise

and Wolf (2001). For three qubits a complete set of inequalities has been found numerically by Pitowsky and Svozil (2000). See also Pitowsky (1989) and Peres (1999).

⁵For an extensive classification of the inequalities see Werner

term to the correlation function is zero. In this way we obtain a local realistic model of a certain correlation function. However, one should check that this construction indeed reproduces the model for the correlation function for the set of settings that enter inequality (15), that is for $E(\vec{n}_{k_1}, \dots, \vec{n}_{k_N})$. For simplicity take $N = 2$. Notice that the expansion coefficients of the four-dimensional vector $(E(\vec{n}_1, \vec{n}_1), E(\vec{n}_1, \vec{n}_2), E(\vec{n}_2, \vec{n}_1), E(\vec{n}_2, \vec{n}_2))$ in terms of orthogonal basis vectors $(s_1 s_2, s_1^2 s_2, s_1 s_2^2, s_1^2 s_2^2)$ (recall that $s_1, s_2 \in \{-1, 1\}$) are equal to the expressions within the moduli entering inequality (15). Next notice, that by the construction shown above the local realistic model for $N = 2$ gives $(E_{LV}(\vec{n}_1, \vec{n}_1), E_{LV}(\vec{n}_1, \vec{n}_2), E_{LV}(\vec{n}_2, \vec{n}_1), E_{LV}(\vec{n}_2, \vec{n}_2)) = \frac{1}{4} \sum_{s_1, s_2} \sum_{k_1, k_2} s_1^{k_1} s_2^{k_2} E(\vec{n}_{k_1}, \vec{n}_{k_2}) (s_1 s_2, s_1^2 s_2, s_1 s_2^2, s_1^2 s_2^2)$. Thus, since the vector built out of the correlation function values and its local realistic counterpart have the same expansion coefficients, they are equal and the sufficiency of (15) as a condition for local realism is proven. The generalization to an arbitrary N is obvious.

Above we derived the full set of Bell inequalities for multi-qubit correlations. This strictly defines the boundary of the validity of local realism. We will now discuss states which violate such inequalities with the specific aim of investigating the way information can be distributed between the subsystems of such states.

VI. N QUBITS THAT VIOLATE LOCAL REALISM

Let us consider the general N -qubit state as in Eq. (10). Then the N -qubit quantum correlation function for a Bell-GHZ type experiment is

$$E_{QM}(\vec{n}_{k_1}, \dots, \vec{n}_{k_N}) = \text{Tr}[\rho(\vec{\sigma} \cdot \vec{n}_{k_1} \otimes \dots \otimes \vec{\sigma} \cdot \vec{n}_{k_N})] \quad (19)$$

$$= \sum_{x_1, \dots, x_N} T_{x_1 \dots x_N} (\vec{n}_{k_1})_{x_1} \dots (\vec{n}_{k_N})_{x_N} \quad (20)$$

where $(\vec{n}_{k_j})_{x_j}$ ($x_j = 1, 2, 3$) are the three Cartesian components of the vector $\vec{n}_{k_j}$. Equation (19) means that the N -particle correlation function is fully defined by a tensor $\hat{T}$ (the indices of which can take values 1, 2, 3, and which belongs to R^{3N}). For convenience we shall write down the last equation in a more compact way as $E_{QM}(\vec{n}_{k_1}, \dots, \vec{n}_{k_N}) = \langle \hat{T}, \vec{n}_{k_1} \otimes \dots \otimes \vec{n}_{k_N} \rangle$, where $\langle \dots, \dots \rangle$ denotes the scalar product in R^{3N} .

The necessary and sufficient condition (15) for a local realistic description of N -particle correlations implies that the quantum correlations for N qubits can always have a local and realistic model for the Bell-type experiment if and only if

$$\sum_{\substack{s_1, \dots, s_N \\ = -1, 1}} |\langle \hat{T}, \sum_{k_1=1}^2 s_1^{k_1} \vec{n}_{k_1} \otimes \dots \otimes \sum_{k_N=1}^2 s_N^{k_N} \vec{n}_{k_N} \rangle| \leq 2^N \quad (21)$$

for any possible choice $\vec{n}_{k_1}, \dots, \vec{n}_{k_N}$ of each observer's two local settings $\vec{n}_1$ and $\vec{n}_2$.

This condition can be simplified further, provided one notices that for each observer there always exist two mutually orthogonal unit vectors $\vec{a}_1$ and $\vec{a}_2$, independently defined for each observer, and the angle α_j such that $\sum_{k_j=1}^2 \vec{n}_{k_j} = 2\vec{a}_1 \cos(\alpha_j + \frac{\pi}{2})$ and $\sum_{k_j=1}^2 (-1)^{k_j} \vec{n}_{k_j} = 2\vec{a}_2 \cos(\alpha_j + \pi)$. Denoting with $c_{x_j} = \cos(\alpha_j + x_j \frac{\pi}{2})$ one can write the inequality (21) as

$$\sum_{\substack{x_1, \dots, x_N \\ = 1, 2}} |c_{x_1} \dots c_{x_N} \langle \hat{T}, \vec{a}_{x_1} \otimes \dots \otimes \vec{a}_{x_N} \rangle| \leq 1. \quad (22)$$

One can rewrite this inequality as

$$\sum_{\substack{x_1, \dots, x_N \\ = 1, 2}} |c_{x_1} \dots c_{x_N} \langle \hat{T}, \vec{a}_{x_1} \otimes \dots \otimes \vec{a}_{x_N} \rangle| \leq 1 \quad (23)$$

where $T_{x_1 \dots x_N}$ is now a component of the tensor $\hat{T}$ in a new set of local coordinate systems, which among their basis vectors have $\vec{a}_1$ and $\vec{a}_2$ which serve as the unite vectors which define the directions x and y .

The necessary and sufficient condition for impossibility of any local realistic description of N -qubit correlations is that the maximum of the left-hand side of inequality (23) is larger than one. Once the values of the elements of the correlation tensor are given for the specific density matrix one can check via maximization procedure whether the local realistic description is possible.

On the other hand, it is easy to notice that with the aid of the Cauchy-Schwarz inequality, one has an inequality

$$\sum_{\substack{x_1, \dots, x_N \\ = 1, 2}} |c_{x_1} \dots c_{x_N} T_{x_1 \dots x_N}| \leq \sqrt{\sum_{\substack{x_1, \dots, x_2 \\ = 1, 2}} T_{x_1 \dots x_2}^2}. \quad (24)$$

Combining inequality (23) with (24) we obtain that a sufficient condition for the possibility of a local realistic description of the quantum N -qubit correlations in any Bell-type experiment is that

$$\sum_{\substack{x_1, \dots, x_N \\ = 1, 2}} T_{x_1 \dots x_N}^2 \leq 1 \quad (25)$$

for all possible choices of local coordinate systems for N qubits as then the full set of Bell inequalities (16) is satisfied. Equivalently, if at least one of the Bell inequalities from the set is violated then condition (25) is violated for at least one choice of local coordinate systems.

Our measure of information (12) is exactly equal to the square of the corresponding element of the quantum correlation matrix. This establishes the equivalence between the condition for local realism (25) on correlation tensor elements and our information-theoretical criterion (7) for information contained in correlations by states which do not reveal quantum entanglement. If the state violates at

least one of the Bell inequalities from the full set (16) then this state is characterized by the information-theoretical criterion (5).

By performing rotations in the x - y planes of the N observers one can vary the values of the elements of the correlation tensor, but these variations do not change the left-hand side of inequality (25). In information-theoretic language we say that the total information content in x - y plane correlations is invariant under these variations. The invariance property implies that one can find local coordinate systems for which some of the correlation tensor elements vanish thus having criterion (25) which involves a smaller number of them (For the case of three qubits see Scarani and Gisin, 2001). For example, in the two-qubit case the rotations in the x - y planes of the two observers are obtained with the use of two parameters, each describing the rotation angle for the given local observer, and therefore one can always find local coordinate systems such that two of correlation tensor elements vanish ($T_{xy} = T_{yx} = 0$) (See Horodecki and Horodecki, 1996). Then it can easily be seen that varying the two angles α_1 and α_2 the expression on the left hand side of inequality (24) can be saturated by the one on the right hand side. Finally this establishes the condition (25) for two qubits as the necessary and sufficient condition for the correlations measured on an arbitrary two-qubit mixed state to be understood within the local realistic picture. In that case our necessary and sufficient condition $I_{xx} + I_{yy} > 1$ (or equivalently $T_{xx}^2 + T_{yy}^2 > 1$) for violation of the most general Bell's inequality is equivalent with the necessary and sufficient condition for violation of the CHSH inequality which was obtained by the Horodeckis (1995). Thus, our result also confirms that non-violation of the CHSH inequality is a necessary and sufficient condition for the local realistic description of two-qubit correlations.

We will now analyze from our information theoretic point the case of an N -qubit Bell-type experiment. We are specifically interested in the limit up to which the experiment still has a local realistic interpretation. Consider the state which is a mixture of the maximally entangled state and the noise induced by experimental imperfections. Such a state is known as the Werner state and has the form

$$\rho_W = V|\psi_{GHZ}\rangle\langle\psi_{GHZ}| + (1 - V)\rho_{noise} \quad (26)$$

where $|\psi_{GHZ}\rangle = \frac{1}{\sqrt{2}}(|+z\rangle_1 \dots |+z\rangle_N + |-z\rangle_1 \dots |-z\rangle_N)$ is the maximal entangled (GHZ) state and $\rho_{noise} = \frac{1}{2^N}I$ is the completely mixed state. Here e.g., $|+z\rangle_j$ denotes the spin up of particle j along z . We would like to emphasize that the weight V of the GHZ-state can operationally be interpreted as the visibility observed in a multi-particle interference experiment (Belinskii and Klyshko, 1993).

For the purposes of our argument we will now calculate the number of non-zero correlation tensor elements

(which are related to our individual measures of information contained in the correlations) for the Werner state. Note first that for any measurement direction $\vec{n}$ belonging to the x - y plane the spin component $\vec{n} \cdot \vec{\sigma}$ has its eigenvectors in the form $|\pm \vec{n}\rangle = \frac{1}{\sqrt{2}}(|+z\rangle \pm e^{i\phi}|-z\rangle)$, where ϕ is the azimuthal angle of the vector $\vec{n}$. Using Eq. (19) one can easily show that the correlation function for arbitrary chosen measurement directions within the local x - y planes is

$$E_W(\phi_1, \dots, \phi_N) = V \cos\left(\sum_{k=1}^N \phi_k\right). \quad (27)$$

This implies that the correlation tensor elements $T_{x_1 \dots x_N}$ with $x_1, \dots, x_N$ each being either x or y are given by $T_{x_1 \dots x_N} = V \cos(m_y \frac{\pi}{2})$, where m_y is the number of y 's in $\{x_1, \dots, x_N\}$. Therefore, for each N one always has $T_{xxx \dots x} = V$. The other components are zero except for those that contain an even number of y 's, which are either V or $-V$. This results in the total number $1 + \sum_{k=1}^{N/2} \binom{N}{2k} = 2^{N-1}$ of non-zero components for the even N and $1 + \sum_{k=1}^{(N-1)/2} \binom{N}{2k} = 2^{N-1}$ for the odd N .

We would like to stress again the equivalence $I_{x_1 \dots x_N} = T_{x_1 \dots x_N}^2$ between the measure of information contained in the correlation between measurements performed along directions $x_1, \dots, x_N$ and the square of the corresponding correlation tensor element. In the case of N qubits in the Werner state we therefore have 2^{N-1} individual measures of information with the value V^2 and we have the remaining ones equal to zero. Inserting these values into the information criterion (7) (or equivalently (25)) one obtains

$$V \leq \left(\frac{1}{\sqrt{2}}\right)^{N-1} \quad (28)$$

for the maximal visibility which still allows the correlations between N qubits in the Werner state to be understood within a local realistic picture. Note that in such a case the right-hand side of the inequality (24) is one. Since the value given on the right-hand side of (28) is also the minimal visibility necessarily to violate the inequalities for N qubits obtained by Belinskii and Klyshko (1993) and since they are included in our set of all possible inequalities (16) we therefore can conclude that our information criterion (7) is the necessary and sufficient condition for correlations between N qubits in the Werner state to violate the local realistic description.

It is interesting to note that different inequalities can be obtained for correlations to be understood within local realism where not only two but three (Żukowski and Kaszlikowski, 1997) or even all possible measurement settings (Żukowski, 1993) are chosen by N observers. There even lower thresholds for the visibilities were obtained to violate the local realistic description. For such an experimental situation our criterion (7) is the sufficient condition for violation of the inequalities.

Another interesting observation is that for N maximally entangled qubits N bits of information rest in the correlations, as opposed to always not more than one bit for the classically composed ones. In the case of GHZ (for $V = 1$ in the consideration given above) the information criterion (7) results in $2^{N-1} \leq 1$ which clearly shows that with growing N the discrepancy between quantum and classical correlations grows exponentially. This is in concurrence with the fact that the GHZ theorem is stronger than Bell's and its strength, as measured by the magnitude of violation of (16) for maximally entangled states, exponentially increases with the number of qubits (Mermin, 1990; Ardehali, 1992, Belinskii and Klyshko, 1993).

VII. CONCLUSIONS

We now would like to review what we have done in the present paper and put it in a broad perspective. The paper contains two independent main approaches to the question of quantum entanglement, and we finally show their essential equivalence.

In the first approach we start from the conceptual position that quantum mechanics is about information. We express the information contained in composite systems such that it can be divided into the information carried by the individuals versus the information contained in the correlations between observations made on the individuals. We further assume that the information contained in any system, be it individual or composite, is finite.

Considering first the classically composed systems we note that any correlations we might observe between the subsystems of such a composite system can simply be understood on the basis of correlations between the properties the individual subsystems have on their own. This means that if we know all properties of the individual subsystems we can definitely conclude how much information is contained in their correlations. For the quantum entangled systems this is not true anymore. Such composite quantum systems can carry more information in joint properties than what may be concluded from knowledge of the individuals. These considerations lead to a natural information-based understanding of quantum entanglement. Within this view we see Mermin's (1998) "correlations without correlata" as reflecting that when correlations are defined there is no information left to define "correlata" as well. "Correlations have physical meaning; that which they correlate does not," as stated by Mermin (1998).

In an independent approach we obtain the most general set of Bell inequalities for N qubits. That way we arrive at a necessary and sufficient condition for quantum states whose correlations cannot be understood within local realism. Local realism is based on the assumption that results of the observations on the individual

systems are predetermined and independent of whatever measurements might be performed distantly. One may notice that this assumption implicitly says that correlations between subsystems do not go beyond what might be concluded from the properties of individual subsystems.

We finally show that the two approaches, the information theoretical one and the one via Bell's inequalities, are equivalent in their essence. This is done via the fact that the Bell inequality criteria can be translated into a statement about correlations (probabilities), which again can be understood as an information theoretical expression. This requires the use of a new measure of information introduced earlier (Brukner and Zeilinger, 1999). This measure of information is distinct from Shannon's measure (Shannon, 1948). The main conceptual difference is that Shannon's measure tacitly assumes that the properties of the systems carrying the information are already well defined prior to, and independent of, observation (Brukner and Zeilinger 2001). In quantum mechanics this clearly is not the case. There, the criterion for choosing the new measure of information was that it is invariant on the experimentalist's free choice of a complete set of mutually complementary observables.

Summing up, we would like to draw the reader's attention to the fact of the equivalence of the two approaches in the present paper, the information theoretic one and the one via Bell's inequalities. It is evident that the first one is both conceptually and formally much simpler. It is suggestive that this new information theoretic formulation of quantum phenomena opens up the avenue of new approaches to well known problems in quantum information physics and in the foundations of quantum mechanics.

ACKNOWLEDGEMENTS

The work was supported by the Austrian Science Foundation (FWF), project F1506, and by the QIPC Program of the European Union. It is also a part of the Austrian-Polish collaboration programme 2400. M.Ž. acknowledges KBN grant No. 5 P03B 088 20.

REFERENCES

- Ardehali, M., 1992, "Bell inequalities with a magnitude of violation that grows exponentially with the number of particles," *Phys. Rev. A* **46**, 5375-5378.
- Aspect, A., P. Grangier, and G. Roger, 1981, "Experimental tests of realistic local theories via Bell's theorem," *Phys. Rev. Lett.* **47**, 460-463.
- Belinskii, A. V. and D. N. Klyshko, 1993, *Phys. Usp.* **36**, 653 (1993).

- Bell, J. S., 1964, "On the Einstein-Podolsky-Rosen paradox," *Physics* **1**, 195-200; reprinted Bell, J. S., 1987, *Speakable and Unspeakable in Quantum Mechanics* (Cambridge Univ. Press, Cambridge).
- Bohr, N., 1958, *Atomic Physics and Human Knowledge* (Wiley, New York).
- Bruckner, Č., and A. Zeilinger, 1999, "Operationally invariant information in quantum measurements," *Phys. Rev. Lett.* **83**, 3354-3357.
- Bruckner, Č. and A. Zeilinger, 2001, "Conceptual inadequacy of the Shannon information in quantum measurements," *Phys. Rev. A* **63**, 022113 1-10.
- Clauser, J., M. Horne, A. Shimony, and R. Holt, 1969, "Proposed experiment to test local hidden-variable theories," *Phys. Rev. Lett.* **23**, 880-884.
- Einstein, A., B. Podolsky, and N. Rosen, 1935, "Can quantum-mechanical description of physical reality be considered complete?," *Phys. Rev.* **47**, 777-780.
- Freedman, S. J., and J. S. Clauser, 1972, "Experimental test of local hidden-variable theories," *Phys. Rev. Lett.* **28**, 938-941.
- Gisin, N., 1991, "Bell's inequality holds for all non-product states," *Phys. Lett. A* **154**, 201-202.
- Gisin, N. and A. Peres, 1992, "Maximal violation of Bell's inequality for arbitrary large spin," *Phys. Lett. A* **162**, 15-17.
- Gnedenko, B. V., 1976, *The Theory of Probability* (Mir Publishers, Moscow).
- Greenberger, D. M., M. Horne, and A. Zeilinger, 1989, "Going beyond Bell's theorem" in *Bell's Theorem, Quantum Theory, and Conceptions of the Universe* edited by M. Kafatos (Kluwer Academic, Dordrecht), p. 73-76.
- Greenberger, D. M., M. Horne, A. Shimony, and A. Zeilinger, 1990, "Bell's theorem without inequalities," *Am. J. Phys.* **58**, 1131-1143.
- Horodecki, R., P. Horodecki, and M. Horodecki, 1995, "Violating Bell inequality by mixed spin-1/2 states: Necessary and sufficient condition," *Phys. Lett. A* **200**, 340-344.
- Horodecki, R. and M. Horodecki, 1996, "Information-theoretic aspects of inseparability of mixed states," *Phys. Rev. A* **54**, 1838-1843.
- Kaszlikowski, D., P. Gnaciski, M. Żukowski, W. Miklaszewski, and A. Zeilinger, 2000, "Violations of Local Realism by Two Entangled N-Dimensional Systems Are Stronger than for Two Qubits," *Phys. Rev. Lett.* **85**, 4418-4421.
- Mermin, N. D., 1990, "Extreme quantum entanglement in a superposition of macroscopically distinct states," *Phys. Rev. Lett.* **65**, 1838-1841.
- Mermin, N. D., 1998, "What is quantum mechanics trying to tell us?" *Am. J. Phys.* **66**, 753-767.
- Pan, J. W., D. Bouwmeester, H. Weinfurter, and A. Zeilinger, 2000, "Experimental test of quantum nonlocality in three photon Greenberger-Horne-Zeilinger entanglement," *Nature* **403**, 515-518.
- Peres A., 1999, "All the Bell inequalities," *Found. Phys.* **29**, 589-614.
- Pitowsky I., 1989, *Quantum Probability - Quantum Logic* (Springer, Berlin).
- Pitowsky I. and K. Svozil, 2001, "Optimal tests of quantum nonlocality," *Phys. Rev. A* **64**, 014102.
- Popescu, S., 1995, "Bell's inequalities and density matrices: Revealing 'hidden' nonlocality," *Phys. Rev. Lett.* **74**, 2619-2622.
- Scarani, V. and N. Gisin, 2001, "Spectral decomposition of Bell's operators for qubits," Preprint quant-ph/0104016 at (<http://xxx.lanl.gov>).
- Schrödinger, E., 1935a, "Die gegenwärtige Situation in der Quantenmechanik," *Naturwissenschaften* **23**, 807-812; 823-828; 844-849. Translation published in *Proc. Am. Phil. Soc.* **124**, p. 323-338 and in *Quantum Theory and Measurement* edited by J. A. Wheeler and W. H. Zurek (Princeton University Press, Princeton), p. 152-167. A copy can be found at (www.emr.hibu.no/lars/eng/cat).
- Schrödinger, E., 1935b, "Discussion of probability relations between separated systems" in *Proceedings of the Cambridge Philosophical Society* **31**, p. 555.
- Shannon, C. E., 1948, "A mathematical theory of communication," *Bell Syst. Tech. J.* **27**, 379. A copy can be found at (<http://cm.bell-labs.com/cm/ms/what/shannonday/paper.html>).
- von Weizsäcker, C. F., 1975, in *Quantum Theory and the Structure of Time and Space II* edited by L. Castell, M. Drieschner and C. F. Weizsäcker (Hanser, München).
- Weih, G., T. Jennewein, C. Simon, H. Weinfurter, and A. Zeilinger, 1998, "Violation of Bell's inequality under strict Einstein locality conditions," *Phys. Rev. Lett.* **81**, 5039-5043.
- Weinfurter, H. and M. Żukowski M., 2001, "Four-Photon Entanglement from Down-Conversion," *Phys. Rev. A* **64**, 010102(R).
- Werner, R. F. and M. M. Wolf, 2001, "All multipartite Bell correlation inequalities for two dichotomic observables per site," Preprint quant-ph/0102024 at (<http://xxx.lanl.gov>).
- Zeilinger, A., 1997, "Quantum teleportation and the non-locality of information," *Phil. Trans. Roy. Soc. Lond.* **1733**, 2401-2404.
- Zeilinger, A., 1999, "A foundational principle for quantum mechanics," *Found. Phys.* **29**, 631-643.
- Żukowski, M., 1993, "Bell Theorem Involving all Settings of Measuring Apparata," *Phys. Lett. A* **177** 290-294.
- Żukowski, M. and Č Bruckner, 2000, "Bell's theorem for general N-qubit states," Preprint quant-ph/0102039 at (<http://xxx.lanl.gov>).
- Żukowski, M. and D. Kaszlikowski, 1997, "Critical visibility for N-particle Greenberger-Horne-Zeilinger correlations to violate local realism," *Phys. Rev. A* **56**, R1682-R1685.